

JOURNAL OF FUNCTIONAL ANALYSIS 47, 143–152 (1982)

A Generalization of Ito's Formula

A. S. ÜSTÜNEL

2, Bd. Auguste Blanqui, Paris 75013, France

Communicated by L. Gross

Received June 12, 1980

INTRODUCTION

In [1], Ito's formula for the Brownian motion is extended to the functions whose Laplacian (in the sense of the distributions) is a measure (cf. [4] for a more probabilistic version). This case seems to be the most general one which can be achieved by the finite dimensional methods. A further generalization of these results would be the case in which the functions are replaced by the distributions; however, evaluation of a distribution at a point of the Euclidean space on which it is defined is meaningless, but there is a concept in the distribution theory which is analogous to the evaluation of the functions at a point, that is, the evaluation of the convolution of a distribution with the Dirac measure corresponding to that point at any \mathcal{C}_0^∞ -function and this is the main idea which has been to the origin of this work.

We start by treating the simplest case, i.e., $B = (B_t; t \geq 0)$ is a d -dimensional standard Brownian motion ($d \geq 1$) and the distributions belong to $\mathcal{S}'(\mathbb{R}^d)$, i.e., the space of the tempered distributions on \mathbb{R}^d . With the help of the Ito's formula, we calculate explicitly $T * \delta_{B_t}$, where $T \in \mathcal{S}'(\mathbb{R}^d)$, δ_{B_t} is the Dirac measure concentrated at B_t and $*$ represents the convolution, as a sum of a local martingale and a process of finite variation with values in $\mathcal{S}'(\mathbb{R}^d)$ (cf. [9, 10, 12]). The second section covers the proof of the existence and the uniqueness of the solutions of the heat equation in $\mathcal{S}'(\mathbb{R}^d)$, and Section III, the results of the first section are extended to the arbitrary \mathbb{R}^d -valued semimartingales and to the elements of $\mathcal{D}'(\mathbb{R}^d)$, i.e., the space of the distributions on \mathbb{R}^d .

I. A GENERALIZATION OF ITO'S FORMULA

We represent by (Ω, \mathcal{F}, P) a completed probability space and by $(\mathcal{F}_t; t \in \mathbb{R}_+)$ an increasing, right continuous filtration of \mathcal{F} such that \mathcal{F}_0 contains all the P -negligible subsets of Ω . If $B = (B_t; t \in \mathbb{R}_+)$ is a standard

d -dimensional Brownian motion, δ_{B_t} defines an $\mathcal{S}'(\mathbb{R}^d)$ -valued semimartingale (cf. [9, 12]). Since as a distribution, $\delta_{B_t(\omega)}$ is of compact support, the convolution $T * \delta_{B_t(\omega)}$ is a well-defined distribution for any $(t, \omega) \in \mathbb{R}_+ \times \Omega$ and it can be expressed as

$$(T * \delta_{B_t})(\phi) = (T(\xi) \otimes \delta_{B_t}(\eta))(\phi(\xi + \eta)) = T(\phi(\cdot + B_t)),$$

where ξ and η denotes the space parameters in \mathbb{R}^d with respect to which the distributions T and δ_{B_t} are defined (cf. [7]) and $\phi \in \mathcal{S}'(\mathbb{R}^d)$. Now we announce:

THEOREM 1.1. *Let $T \in \mathcal{S}'(\mathbb{R}^d)$, then one has the following relation:*

$$T * \delta_{B_t} = T - \int_0^t \nabla T * \delta_{B_s} \cdot dB_s + \frac{1}{2} \int_0^t \Delta T * \delta_{B_s} ds$$

where $\nabla T * \delta_{B_t}$ denotes $(\partial T * \delta_{B_t} / \partial x_1, \dots, \partial T * \delta_{B_t} / \partial x_d)$ and the stochastic integral is defined as

$$\left(\int_0^t \nabla T * \delta_{B_s} \cdot dB_s \right) (\phi) = - \sum_{i=1}^d \int_0^t T * \delta_{B_s} \left(\frac{\partial \phi}{\partial x_i} \right) dB_s^i$$

for $B_t = (B_t^1, \dots, B_t^d)$.

Remark. Since the mapping $x \rightarrow \tau_x \phi$ defined as $\tau_x \phi(y) = \phi(x + y)$ for $\phi \in \mathcal{S}'(\mathbb{R}^d)$ is continuous with values in $\mathcal{S}'(\mathbb{R}^d)$, the process $T * \delta_{B_t}$ has weakly continuous trajectories and $\mathcal{S}'(\mathbb{R}^d)$ being a \mathcal{DS} -space, $T * \delta_{B_t}$ is locally bounded (cf. [9]), hence the stochastic integral is well defined.

We shall prove the theorem in several steps. For the sake of simplicity we suppose $d = 1$. First, if $\phi \in \mathcal{S}'(\mathbb{R})$, by Ito's formula (cf. [3]) we have

$$\phi(\xi + B_t) = \phi(\xi) + \int_0^t \phi'(\xi + B_s) dB_s + \frac{1}{2} \int_0^t \Delta \phi(\xi + B_s) ds.$$

The mapping defined by $(t, \omega) \rightarrow \phi(\cdot + B_t(\omega))$ is a stochastic process with values in $\mathcal{S}'(\mathbb{R})$, hence $\phi(\xi + B_t) - \phi(\xi)$ has also the same property. Moreover $\xi \rightarrow \frac{1}{2} \int_0^t \Delta \phi(\xi + B_s) ds$ belongs also to $\mathcal{S}'(\mathbb{R})$ since

$$\left| \xi^k D_t^p \int_0^t \psi(\xi + B_s) ds \right| \leq |\xi|^k \int_0^t |D_t^p \psi(\xi + B_s)| ds \xrightarrow{|\xi| \rightarrow \infty} 0 \quad \text{a.e.}$$

for any $p, k \in \mathbb{N}$. Consequently $(t, \omega) \rightarrow \int_0^t \phi'(\cdot + B_s) dB_s$ is a stochastic process with values in $\mathcal{S}'(\mathbb{R})$, and we can write

$$(T * \delta_{B_t})(\phi) = T(\phi) + T \left(\int_0^t \phi'(\cdot + B_s) dB_s \right) + \frac{1}{2} T \left(\int_0^t \Delta \phi(\cdot + B_s) ds \right).$$

If we can show that $T \in \mathcal{S}'(\mathbb{R})$ commutes with the integrals, the theorem would be proved. From now on $\mathcal{S}(\mathbb{R})$ and $\mathcal{S}'(\mathbb{R})$ will be denoted respectively by \mathcal{S} and \mathcal{S}' .

LEMMA I.1. *For any $T \in \mathcal{S}'$ and $\psi \in \mathcal{S}$, one has the following relation:*

$$T \left(\int_0^t \psi(\cdot + B_s) ds \right) = \int_0^t T(\psi(\cdot + B_s)) ds \quad \text{a.e. for any } t \geq 0.$$

Proof. Define T_n as $T_n = \inf\{t: |B_t| > n\}$, then T_n is an \mathcal{F}_t -stopping time and $T_n \uparrow + \infty$ as n tends to infinity. If $\psi \in \mathcal{S}$, then the process $(t, \omega) \rightarrow \psi(\cdot + B_{t \wedge T_n}(\omega)) \equiv \psi(\cdot + B_t^n(\omega))$ is bounded. Let K_n be the closed absolutely convex hull in \mathcal{S} of the set

$$\{\psi(\cdot + B_t^n(\omega)) \in \mathcal{S}: (t, \omega) \in \mathbb{R}_+ \times \Omega\}.$$

Since \mathcal{S} is a Montel space, K_n is compact in \mathcal{S} ; denote by $\mathcal{S}[K_n]$ the completion of the subspace spanned by K_n with respect to the norm $g_{K_n}(\phi) = \inf\{\lambda > 0: \phi \in \lambda K_n\}$. Let K_n^0 be the polar of K_n in \mathcal{S}' , and denote by $\mathcal{S}'(K_n^0)$ the completion of the quotient space $\mathcal{S}'/r_{K_n^0}^{-1}(0)$ with respect to the seminorm $r_{K_n^0}$ which is the gauge function of K_n^0 , i.e., if $F \in \mathcal{S}'$, then $r_{K_n^0}(F) = \inf\{\lambda > 0: F \in \lambda K_n^0\}$. It is well known that $(\mathcal{S}[K_n], \mathcal{S}'(K_n^0))$ forms a dual pair (cf. [6]) and if π_n denotes the canonical mapping from \mathcal{S}' onto $\mathcal{S}'(K_n^0)$, then for any $T \in \mathcal{S}'$, $\phi \in \mathcal{S}[K_n]$, one has

$$\langle \phi, \pi_n(T) \rangle = T(\phi),$$

where $\langle \cdot, \cdot \rangle$ is the bilinear form associated to the dual pair $(\mathcal{S}[K_n], \mathcal{S}'(K_n^0))$.

The stochastic process $(t, \omega) \rightarrow \psi(\cdot + B_t^n(\omega))$ is weakly continuous, adapted with values in $\mathcal{S}[K_n^0]$, moreover it is bounded. Hence the Lebesgue integral converges in $\mathcal{S}[K_n^0]$. If $T \in \mathcal{S}'$, then

$$\begin{aligned} \left\langle \pi_n(T), \int_0^t \psi(\cdot + B_s^n) ds \right\rangle &= \int_0^t \langle \pi_n(T), \psi(\cdot + B_s^n) \rangle ds \\ &= \int_0^t T(\psi(\cdot + B_s^n)) ds. \end{aligned}$$

Moreover, in [9], we have showed that this result is independent of the choice of the compact set K_n absorbing $\psi(\cdot + B^n)$; since $T_n \uparrow + \infty$, the lemma is proved. Q.E.D.

We have the analogous result for the stochastic integral:

LEMMA I.2. For any $\psi \in \mathcal{S}$ and $T \in \mathcal{S}'$, one has the following relation:

$$T \left(\int_0^t \psi(\xi + B_s) dB_s \right) = \int_0^t T(\psi(\cdot + B_s)) dB_s \quad \text{a.e. for any } t \geq 0.$$

Proof. With the notation of the proof of Lemma I.1, we have

$$\begin{aligned} T \left(\int_0^t \psi(\xi + B_s^n) dB_s \right) &= \left\langle \pi_n(T), \int_0^t \psi(\cdot + B_s^n) dB_s \right\rangle \\ &= \int_0^t T(\psi(\cdot + B_s^n)) dB_s \end{aligned}$$

since $\psi(\cdot + B^n)$ is bounded, predictable with values in $\mathcal{S}[K_n]$. Moreover, as it is shown in [9, Theorem II.4], this result is independent of the choice of K_n . Q.E.D.

Now, the proof of Theorem I.1 is easy to see by writing

$$\begin{aligned} (T * \delta_{B_t})(\phi) &= T(\phi) + \int_0^t T \left(\frac{\partial \phi}{\partial x}(\cdot + B_s) \right) dB_s + \frac{1}{2} \int_0^t T \left(\frac{\partial^2 \phi}{\partial x^2}(\cdot + B_s) \right) ds \\ &= T(\phi) - \int_0^t \frac{\partial T * \delta_{B_s}}{\partial x}(\phi) dB_s + \frac{1}{2} \int_0^t \frac{\partial^2 T * \delta_{B_s}}{\partial x^2}(\phi) ds. \end{aligned}$$

II. HEAT EQUATION

In this section using Theorem I.1 we study the heat equation in \mathcal{S}' . First let $T \in \mathcal{S}'$ such that

$$E \int_0^t |T * \delta_{B_s}(\phi)|^2 ds < +\infty \quad \text{for } t \geq 0, \phi \in \mathcal{S}.$$

Then $\int_0^t T * \delta_{B_s} dB_s$ is a martingale with values in \mathcal{S}' . Taking the expectation of $T * \delta_{B_t}$ we find, for any $\phi \in \mathcal{S}$, that

$$\begin{aligned} H_t(\phi) &\equiv E[T * \delta_{B_t}(\phi)] = T(\phi) + \frac{1}{2} \int_0^t E[\Delta T * \delta_{B_s}(\phi)] ds \\ &= T(\phi) + \frac{1}{2} \int_0^t E[T * \delta_{B_s}(\Delta \phi)] ds \\ &= T(\phi) + \frac{1}{2} \int_0^t \Delta H_s(\phi) ds, \end{aligned}$$

hence H_t , defined as $E[T * \delta_{B_t}(\phi)] = H_t(\phi)$, is a solution of the heat equation

$$\frac{\partial U}{\partial t} = \frac{1}{2} \Delta U, \quad U(0) = T \in \mathcal{S}'.$$

In fact, using an extension of Ito's formula we will prove a uniqueness result. To do so, we need the following result:

LEMMA II.1. *Suppose that $H: \mathbb{R} \rightarrow \mathcal{S}'(\mathbb{R}^d)$ is a continuous mapping of class \mathcal{C}^1 , i.e., for any $\phi \in \mathcal{S}(\mathbb{R}^d)$, the function $t \rightarrow H_t(\phi)$ is of class \mathcal{C}^1 . Then one has the following relation:*

$$H_t * \delta_{B_t} = H_0 - \int_0^t \nabla_x H_s * \delta_{B_s} \cdot dB_s + \int_0^t \left[\frac{dH_s}{ds} * \delta_{B_s} + \frac{1}{2} \Delta_x H_s * \delta_{B_s} \right] ds. \quad (\text{II.1})$$

Proof. Let T be a positive number and denote by \mathcal{X} the completion of $\mathcal{C}_0^\infty((0, T))$, i.e., the infinitely differentiable functions on $[0, T]$, whose supports are in $(0, T)$, with respect to the seminorm defined as

$$f \rightarrow \|f\|^2 = \int_0^T |f(t)|^2 dt + \int_0^T \left| \frac{df(t)}{dt} \right|^2 dt.$$

Then, the closed graph theorem implies that the mapping $\phi \rightarrow H(\phi)$ is continuous on $\mathcal{S}(\mathbb{R}^d)$ with values in \mathcal{X} , so it is a nuclear mapping. Hence it can be represented as (cf. [6])

$$H_t = \sum_{i=1}^{\infty} \lambda_i F^i \otimes f_i(t) \quad dt\text{-a.e.},$$

where $(\lambda_i) \in l^1$, $\{f_i\}$ is a bounded sequence in \mathcal{X} and $\{F^i\}$ is equicontinuous in $\mathcal{S}'(\mathbb{R}^d)$. By Theorem I.1, $H_t * \delta_{B_t}$ can be written in the differential form as

$$\begin{aligned} d(H_t * \delta_{B_t}) &= \sum_i \lambda_i [F^i * \delta_{B_t} df_i(t) + f_i(t) d(F^i * \delta_{B_t})] \\ &= \sum_i \lambda_i \left[F^i * \delta_{B_t} \frac{df_i}{dt} dt - f_i(t) \nabla_x F^i * \delta_{B_t} \cdot dB_t \right. \\ &\quad \left. + \frac{1}{2} f_i(t) \Delta_x F^i * \delta_{B_t} dt \right] \\ &= - \sum_i \lambda_i f_i(t) \nabla_x F^i * \delta_{B_t} \cdot dB_t + \sum_i \lambda_i \frac{df_i(t)}{dt} F^i * \delta_{B_t} dt \\ &\quad + \frac{1}{2} \sum_i \lambda_i f_i(t) \Delta_x F^i * \delta_{B_t} dt. \end{aligned}$$

Hence for an arbitrary representation of H we have

$$\begin{aligned} H_t * \delta_{B_t} &= H_0 - \sum_i \lambda_i \int_0^t f_i(s) \nabla_x F^i * \delta_{B_s} \cdot dB_s + \sum_i \lambda_i \int_0^t \frac{df_i(s)}{ds} F^i * \delta_{B_s} ds \\ &\quad + \frac{1}{2} \sum_i \lambda_i \int_0^t f_i(s) \Delta_x F^i * \delta_{B_s} ds. \end{aligned}$$

The proof will be completed if we show that the sum commutes with the integrals. In order to see this, stop (B_t) as in the proof of Lemma I.1. Then it is not difficult to see that the integral

$$\int_0^{t \wedge T_n} \sum_i \lambda_i \nabla_x F^i * \delta_{B_s} f_i(s) \cdot dB_s$$

converges in $\mathcal{S}'(\mathbb{R}^d) \hat{\otimes} L^2(\Omega, \mathcal{F}, P)$, i.e., the completed projective tensor product of $\mathcal{S}'(\mathbb{R}^d)$ and $L^2(\Omega, \mathcal{F}, P)$. For the other integrals, the proof is similar and for any $\phi \in \mathcal{S}(\mathbb{R}^d)$ we have

$$\begin{aligned} H_t * \delta_{B_t}(\phi) &= H_0(\phi) - \int_0^t \nabla_x H_s * \delta_{B_s}(\phi) \cdot dB_s \\ &\quad + \int_0^t \left[\frac{dH_s}{ds} * \delta_{B_s}(\phi) + \frac{1}{2} \Delta_x H_s * \delta_{B_s}(\phi) \right] ds \end{aligned}$$

$dP \times dt$ — a.e., but the right-hand side is continuous in t hence this relation is true up to an evanescent process; since T is arbitrary, the proof is completed.

Q.E.D.

Remark. Instead of \mathcal{X} one can choose the completion of $\mathcal{C}_0^\infty((0, T))$ with respect to the seminorm

$$f \rightarrow \sup_{0 \leq t \leq T} \left[|f(t)| + \left| \frac{df(t)}{dt} \right| \right]$$

and with this choice the representation of H_t , holds for all t and the proof becomes simpler.

In order to prove the uniqueness of the solution of the heat equation, we shall use the following result:

LEMMA II.2. *Let $T \in \mathcal{S}'(\mathbb{R}^d)$ be such that*

$$E \left[\int_0^t (T * \delta_{B_s}(\phi))^2 ds \right] < +\infty \quad \text{for any } t \in \mathbb{R}_+, \phi \in \mathcal{S}(\mathbb{R}^d),$$

then for any $\phi \in \mathcal{S}(\mathbb{R}^d)$

$$E \left[\int_0^t (H_{a-s} * \delta_{B_s}(\phi))^2 ds \right] < +\infty \quad \text{for any } t \leq a, \text{ da-a.e.,}$$

where H_a is defined as $H_a(\phi) = E[T * \delta_{B_a}(\phi)]$.

Proof. Without loss of generality, we may suppose that (B_t) is a Markov process with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. Moreover, we can take T in the form $D_t^\beta g$ where g is a continuous polynomially bounded function (cf. [5, p. 139]) and D_t^β represents the differential operator $D_{t_1}^{\beta_1} \dots D_{t_d}^{\beta_d}$ with $\beta_1 + \dots + \beta_d = \beta$ and $D_{t_i}^{\beta_i} = \partial^{\beta_i} / \partial \xi_i^{\beta_i}$. Then, if $\phi \in \mathcal{S}(\mathbb{R}^d)$, we have

$$\begin{aligned} & \int_0^t (H_{a-s} * \delta_{B_s}(\phi))^2 ds \\ &= \int_0^t \left| \int_{\mathbb{R}^d} g(\xi) E_{\tilde{\omega}}[D_t^\beta \phi(\xi + B_{a-s}(\tilde{\omega}) + B_s(\omega))] d\xi \right|^2 ds, \quad (\text{II.2}) \end{aligned}$$

where $E_{\tilde{\omega}}$ represents the mathematical expectation with respect to $\tilde{\omega}$. If we denote by E_x the mathematical expectation with respect to Wiener measure W with $W\{B_0 = x\} = 1$, (II.2) can be written

$$\begin{aligned} & \int_0^t \left| \int_{\mathbb{R}^d} g(\xi) E_{B_s(\omega)}[D_t^\beta \phi(\xi + B_{a-s})] d\xi \right|^2 ds \\ &= \int_0^t \left| \int_{\mathbb{R}^d} g(\xi) E_0[D_t^\beta \phi(\xi + B_a) | \mathcal{F}_s] d\xi \right|^2 ds \\ &\leq \int_0^t E_0 \left[\left| \int_{\mathbb{R}^d} g(\xi) D_t^\beta \phi(\xi + B_a) d\xi \right|^2 \middle| \mathcal{F}_s \right] ds \\ &= \int_0^t E_0[|T * \delta_{B_a}(\phi)|^2 | \mathcal{F}_s] ds. \end{aligned}$$

Consequently one has

$$E \int_0^t (H_{a-s} * \delta_{B_s}(\phi))^2 ds \leq t E[|T * \delta_{B_a}(\phi)|^2] < +\infty \quad \text{da-a.e. Q.E.D.}$$

Now the uniqueness can be proved:

THEOREM II.1. Suppose that $T \in \mathcal{S}'(\mathbb{R}^d)$ satisfies the following relation:

$$E \int_0^t |T * \delta_{B_s}(\phi)|^2 ds < +\infty$$

for any $\phi \in \mathcal{S}(\mathbb{R}^d)$. Then the mapping $H: R_+ \rightarrow \mathcal{S}'(\mathbb{R}^d)$ defined as $H_t(\phi) = E[T * \delta_{B_t}(\phi)]$ is a solution of

$$\frac{dH}{dt} = \frac{1}{2} \Delta H, \quad t > 0,$$

$$H_0 = T.$$

Moreover, if G is another solution with weakly differentiable trajectories then, $G = H$.

Proof. The existence has been proved at the beginning of this section. The uniqueness goes on as in the finite dimensional case: By Lemma II.1 for any $a > 0$

$$H_{a-t} * \delta_{B_t} = H_a + M_t^a,$$

where M_t^a is a local martingale; but by Lemma II.2, it is da -almost everywhere a martingale. Taking the mathematical expectation we find for $t = a$

$$E[T * \delta_{B_a}] = H_a, \quad da\text{-a.e.}$$

hence $G_a = H_a$ da -a.e.; moreover they have the weakly continuous trajectories and the uniqueness follows. Q.E.D.

Remark. In fact this result can be extended to those G whose trajectories are almost everywhere differentiable.

III. EXTENSIONS TO SEMIMARTINGALES

In this section we shall work again in the one-dimensional cases; however the results can be extended trivially to any finite dimensional Euclidean space.

Let X be a semimartingale. If $\phi \in \mathcal{S}$, then one has (cf. [3])

$$\begin{aligned} \phi(X_t + \xi) &= \phi(X_0 + \xi) + \int_0^t \phi'(X_{s-} + \xi) dX_s \\ &\quad + \frac{1}{2} \int_0^t \phi''(X_{s-} + \xi) d\langle X^c, X^c \rangle_s \\ &\quad + \sum_{0 \leq s < t} [\phi(\xi + X_s) - \phi(\xi + X_{s-}) - \phi'(\xi + X_{s-}) \Delta X_s], \end{aligned}$$

where X_{t-} represents the left limit of X_t at t and $\Delta X_s = X_s - X_{s-}$. We have

LEMMA III.1. *Let*

$$A_t(\omega, \xi) = \sum_{0 \leq s \leq t} [\phi(\xi + X_s) - \phi(\xi + X_{s-}) - \phi'(\xi + X_{s-}) \Delta X_s],$$

then $(t, \omega) \rightarrow A_t(\omega, \cdot)$ is an adapted, right continuous stochastic process with values in \mathcal{S} .

Proof. Let $K_t(\omega) = \sup_{s \leq t} |X_s(\omega)|$. Then $X_s(\omega) \in [-K_t(\omega), K_t(\omega)]$ and $\Delta X_s(\omega) \in [-2K_t(\omega), 2K_t(\omega)]$ for any $s \leq t$. Hence the set

$$M_t(\omega) = \{\phi(\cdot + \Delta X_s(\omega)): 0 \leq s \leq t\}$$

is relatively compact in \mathcal{S} . If q is any continuous seminorm on \mathcal{S} , by the Taylor's formula, we have

$$q(A_t(\omega, \cdot)) \leq \sum_{s \leq t} (\Delta X_s)^2 q'(\phi(\cdot + \theta \Delta X_s(\omega))), \quad \theta \in [-1, 1],$$

where q' is also a continuous seminorm. Hence

$$q(A_t(\omega, \cdot)) \leq \sup\{q'(\phi): \phi \in M_t(\omega)\} \cdot \sum_{s \leq t} (\Delta X_s)^2 < +\infty;$$

consequently $A_t(\omega)$ converges in \mathcal{S} for almost everywhere ω . Measurability follows from the fact that the mapping

$$(t, \omega, \xi) \rightarrow \phi(\xi + X_t(\omega))$$

is measurable since it can be written a

$$((t, \omega), \xi) \rightarrow (X_t(\omega), \xi) \circ y \rightarrow \phi(y). \quad \text{Q.E.D.}$$

THEOREM III.1. *For any tempered distribution T one has the following relation:*

$$\begin{aligned} T * \delta_{X_t} &= T * \delta_{X_0} - \int_0^t T' * \delta_{X_{s-}} dX_s + \frac{1}{2} \int_0^t T'' * \delta_{X_{s-}} d\langle X^c, X^c \rangle_s \\ &\quad + \sum_{0 \leq s \leq t} [T * \delta_{X_s} - T * \delta_{X_{s-}} + T' * \delta_{X_{s-}} \Delta X_s], \end{aligned}$$

where X^c represents the continuous local martingale part of X .

Proof. By Lemma III.1 the term corresponding to the jumps of X is well defined; for the other terms, one proceeds exactly as in the proofs of Theorem I.1 and Lemma I.1. Q.E.D.

Remark. Let us note that this result extends almost trivially to \mathcal{D}' , i.e.,

the space of the distributions on \mathbb{R} (hence to $\mathcal{D}'(\mathbb{R}^d)$). To see this it is sufficient to remark that the mapping

$$x \rightarrow \phi(\cdot + x) = \tau_x \phi$$

is continuous from \mathbb{R} into \mathcal{D} for any $\phi \in \mathcal{D}$; hence the mapping $(t, \omega) \rightarrow \tau_{X_t(\omega)}(\phi)$ is measurable and (with the notations of Lemma III.1) $M_t(\omega) = \{\phi(\cdot + \Delta X_s(\omega)): 0 \leq s \leq t\}$ is a bounded set in \mathcal{D} .

REFERENCES

1. G. A. BROSAMLER, Quadratic variation of potentials and harmonic functions, *Trans. Amer. Math. Soc.* **149** (1970), 243–257.
2. R. M. BLUMENTHAL AND R. K. GETTOOR, "Markov Processes and Potential Theory," Academic Press, New York, 1968.
3. P. A. MEYER, Un cours sur les intégrales stochastiques, in "Séminaire de Probabilités X," Lecture Notes in Mathematics No. 511, Springer-Verlag, Berlin/Heidelberg/New York, 1976.
4. P. A. MEYER, La formule d'Ito pour le mouvement brownien d'après G. Brosamler, in "Séminaire de Probabilités XII," Lecture Notes in Mathematics No. 649, Springer-Verlag, Berlin/Heidelberg/New York, 1978.
5. M. REED AND B. SIMON, "Methods of Modern Mathematical Physics, I. Functional Analysis," Academic Press, New York, 1972.
6. H. H. SCHAEFER, "Topological Vector Spaces," Graduate Texts in Mathematics, Springer-Verlag, Berlin/Heidelberg/New York, 1970.
7. L. SCHWARTZ, "Théorie des distributions," Hermann, Paris, 1966.
8. B. SIMON, "Functional Integration and Quantum Physics," Academic Press, New York/San Francisco/London, 1979.
9. A. S. ÜSTÜNEL, Stochastic integration on nuclear spaces and its applications, *Ann. Inst. H. Poincaré*, in press.
10. A. S. ÜSTÜNEL, A characterization of semimartingales on the nuclear spaces, *Z. Wahrsch. Verw. Gebiete*, in press.
11. A. S. ÜSTÜNEL, Some applications of stochastic integration in infinite dimension, *Stochastics*, in press.
12. A. S. ÜSTÜNEL, Formule de Feynman–Kac stochastique, *C. R. Acad. Sci. Paris* **292** (1981), 595–597.
13. A. S. ÜSTÜNEL, "Calcul stochastique sur les espaces nucléaires et ses applications," Thèse de Doctorat d'Etat, Université de Paris VI, 1981.